

Medical Imaging Consultants of New York, P.C.

240 Madison Avenue 2nd floor New York, N.Y. 10016-2820

Tel: 212 744-6200 • Fax: 212 490-3933

Joseph Tuvia, M.D.

Jane Ebin Tuvia, M.D.

██████████, Elizabeth
January 15, 2013

Page 2:

with a tear. There is a radial tear in the posterior horn of the lateral meniscus. Increase signal intensity seen in the lateral meniscus is consistent with meniscal degeneration. There is irregularity of the patellar articulating surface with subchondral cystic changes, and cartilaginous loss, degenerative in nature. The periarticular soft tissue structures are normal.

Impression:

Posterior horn medial meniscal tear, likely degenerative in nature.

Radial tear. Posterior horn lateral meniscus, likely degenerative as well.

Intrasubstance tear, posterior cruciate ligament, likely chronic.

Degenerative changes about the patellofemoral compartment.

Moderate joint effusion, either inflammatory or posttraumatic. Also seen is a popliteal fossa cyst.

STATEMENT: I declare, under the penalties of perjury, that the information contained within this document was prepared and is the work product of the undersigned, and is true to the best of my knowledge and information.

Sincerely,



Joseph Tuvia, M.D.

Lic. 187294-1