

Medical Imaging Consultants of New York, P.C.

240 Madison Avenue 2nd floor New York, N.Y. 10016-2820

Tel: 212 744-6200 • Fax: 212 490-3933

Joseph Tuvia, M.D.

Jane Ebin Tuvia, M.D.

██████████, Elizabeth
January 15, 2013

Page 2:

the infraspinatus muscle. There is partial tear of the subscapularis tendon. The teres minor tendon is intact. The glenoid labrum and glenohumeral compartment are intact. The long head bicipital tendon is not visualized in its expected location, likely torn and retracted. The short head bicipital tendon is intact.


Impression:

Advanced degenerative changes about the acromioclavicular joint. Full thickness tear of the distal supraspinatus tendon, retraction and atrophy. Partial tear, infraspinatus tendon and subscapularis tendon, mild infraspinatus muscular atrophy. Large joint effusion and fluid in the subacromial subducted bursa. There is a tear of the long head bicipital tendon with tendinous retraction.

The constellation of the above findings is most likely degenerative in nature and long-standing.

STATEMENT: I declare, under the penalties of perjury, that the information contained within this document was prepared and is the work product of the undersigned, and is true to the best of my knowledge and information.

Sincerely,


Joseph Tuvia, M.D.

Lic. 187294-1

15E